

- [Home](#)
- [Editorial Board](#)
- [Submit a paper](#)
- [Indexing and Awards](#)
- [Reviewers](#)
- [Articles in Press](#)
- [Publication Ethics](#)
- [Archives](#)

**JESTEC**  
Journal of Engineering  
Science and Technology



WEB OF SCIENCE™



- **Editor-in-Chief**  
**Assoc Prof Dr Abdulkareem Sh. Mahdi Al-Obai**  
Taylor's University, Malaysia
- [Editorial Board](#)

Journal of Engineering Science and Technology (JESTEC) indexed by [SCOPUS](#) since 2010.

Journal of Engineering Science and Technology has been selected for coverage in [Clarivate Analytics products and services](#).

Beginning with 2016, this publication will be indexed abstracted in:

[Emerging Sources Citation Index \(ESCI\)](#)

**ISSN: 1823-4690**

## Aims and Scope

JESTEC (Journal of Engineering Science and Technology) is a peer-reviewed journal that aims to publish and disseminate original research articles latest developments in all fields of engineering science and technology. The journal publishes original papers in English, which contribute to underest engineering science and improving engineering technology and education. The articles may be theoretical (including computational), experime both. The contribution should be unpublished before and not under consideration for publication elsewhere.

JESTEC maintains a standard double-blind peer-review process. The double-blind process means that the author and the reviewer's ident not known to each other.

JESTEC is an Open Access journal and does not charge readers or their institutions to access the journal articles. Open access supports rights to read, download, copy, distribute, print, search, or link to these articles' full texts, provided they are properly acknowledged and cite

JESTEC publishes six issues per year.

The publication fees in JESTEC include submission, reviewing, editing, publishing, and uploading the accepted article to the JESTEC websit all these services, JESTEC charges USD375 or MYR1550 per paper (inclusive 6% SST). The number of pages per paper should be a minimu pages and a maximum of 15-18 pages.

## Subscriptions and enquiries

Contact the [Executive Editor](#)

## Aims & Scope

[Home](#)

[Editorial Board](#)

[Submit a paper](#)

[Indexing and Awards](#)

[Reviewers](#)

[Articles in Press](#)

[Publication Ethics](#)

[Archi](#)



**Volume 16, Issue 1, February 2021**

**Page 0001 - 0887**

[Contrast enhancement using adaptive threshold based dynamic range adjustment in LUV colour space](#)

P. M. Pardhi, S. D. Thepade

1 - 24

[Economic operation planning of combined heat and power smart distribution system](#)

M. K. Al-Saadi

25 - 43

[The strategic role of smart city \(Bandung care\) in the development of Bandung city](#)

D. S. Soegoto, Y. Pangestika

44 - 51

[X-ray transmission-rate-measuring by using low-cost laboratory counter](#)

A. T. Zengin

52 - 64

[Robust energy efficient control for wireless sensor networks via unity sliding mode](#)

S. M. Raafat, A. M. Mahmood

65 - 84

[Analysis of distributed deep-learning based digital learning media using thin client devices for inclusion vocational school students](#)

A. Andriana, A. Ana, H. Puspita, I. Y. Wulandari

85 - 91

[Grid-independent PV-wind-diesel generator hybrid renewable energy system for a medium population: A case study](#)

Z. W. J. Al-Shammari, M. M. Azizan, A. S. F. Rahman

92 - 106

[Characterization of the earth dam embankment using shallow seismic data: A case study at Klong Sadao dam, southern Thailand](#)

S. Yordkayhun, K. Wattanasen

107 - 121

[Water quality modelling and management of Diyala river and its impact on Tigris River](#)

B. Sh. Abed, M. H. Daham, A. H. Ismail

122 - 135

[Performance of green mortar made from locally available waste tiles and silica fume](#)

M. K. Obaid, M. S. Nasr, I. M. Ali, A. A. Shubbar, K. S. Hashim  
136 - 151

[Using monkey optimization algorithm to detect neris botnet](#)

I. A. Saleh, M. A. Kamal, L. M. Ibrahim  
152 - 164

[Vetiver waste fertilizer as a growth catalyst on mung bean](#)

E. S. Soegoto, R. Jumansyah, S. Luckyardi  
165 - 175

[A review of the transformation techniques in the OFDM system](#)

M. H. M. Nerma  
176 - 193

[Real-time moving objects detection and tracking using deep-stream technology](#)

N. H. Abdulghafoor, H. N. Abdullah  
194 - 208

[Chemotherapy for cervical cancer patients: Is it a "Nightmare"?](#)

Suwendar, A. Fudholi, T. M. Andayani, H. S. Sastramihardja  
209 - 223

[Optimization of adsorption process parameters of Ni \(II\) in a bed packed-column with agro-industrial residue of yam \(dioscorea rotundata\)](#)

C. Tejada-Tovar, A. Villabona-Ortiz, M. Jiménez, A. Herrera-Barros, R. Ortega-Toro  
224 - 235

[Noma multiuser detection with MIMO for maximizing the system capacity and eliminate interference](#)

A. V. Kulkarni, S. N. Motade  
236 - 250

[Learning antenna simulation to enhance technical competency of avionics students](#)

I. Y. Wulandari, E. Ema, A. Ana, N. Indroasyoko, H. Puspita, Andriana, R. Hidayat  
251 - 258

[Preparation of copper aluminum-biochar composite as adsorbent of malachite green in aqueous solution](#)

N. R. Palapa, T. Taher, A. F. Badri, Suheryanto, R. Mohadi, A. Rachmat, A. Lesbani  
259 - 274

[Expressive shares visual secret sharing using discrete cosine transform for improved extracted secret quality](#)

R. Chaturvedi, S. D. Thepade, S. Ahirrao  
275 - 290

[Elicitation of stage discharge relationship for multi-gate controlled barrage](#)

M. S. Shamkhi, N. M. Neama  
291 - 304

[Transfer function modeling and optimization speed response of BLDC Motor E-bike using intelligent controller](#)

I. Anshory, I. Robandi, Jamaaluddin, A. Fudholi, Wirawan  
305 - 324

[Hybridization and modification of the PSO algorithm and its use in personal recognition by OPG X-ray](#)

D. S. Hammadi, A. N. Younis, F. M. Ramo  
325 - 338

[Data visualization for content marketing domain in social media](#)

D. Dharmayanti, A. M. Bachtiar, F. N. Fakhrol

[The use of ultraviolet \(UV\) spectroscopy and chemometrics to quantify the percentages of adulteration in Kalosi ground roasted specialty coffee](#)

D. Suhandy, M. Yulia  
350 - 364

[Low voltage ride through capability enhancement using series connected fact devices in wind energy conversion system](#)

B. P. Ganthia, S. K. Barik, B. Nayak  
365 - 384

[Projection simulation media in technical drawing for vocational students](#)

D. Rohendi, N. Utami, A. Ana, A. Sukandar  
385 - 397

[Static polygon formation in leader- follower robotic system by utilizing RP LIDAR sensor in unknown environment](#)

B. A. Issa, A. T. Rashid  
398 - 414

[Effect of structural defects and impurities on electrical characteristics of photovoltaic cells: Multiscale modelling](#)

O. A. Al-Ani, A. M. A. Sabaawi, E. H. Sadiq, M. F. Mosleh  
415 - 426

[Improvement of cassiterite levels with variation of feed roll and separation roll speed in high tension roll separator \(HTRS\)](#)

L. Pulungan, D. N. Usman, S. Widayati, S. Maulida  
427 - 437

[Assessment of contamination vulnerability of groundwater using susceptibility index method: Semarang city as case study](#)

T. T. Putranto, A. A. Marjuanto  
438 - 454

[Street dust contamination by heavy metals in Babylon governorate, Iraq](#)

O. J. Al-Sareji, R. A. Grmasha, J. M. Salman, K. S. Hashim  
455 - 469

[The use of psychrometric analysis kit for learning air conditioning process in vocational education](#)

E. T. Berman, I. Hamidah, B. Mulyanti, A. Setiawan  
470 - 481

[Feature fusion approach for image retrieval with ordered color means based description of key points extracted using local detectors](#)

J. H. Dewan, S. D. Thepade  
482 - 509

[Characterization of tyre rubber ash and crumb as fine aggregate resource](#)

S. Shahidan, S. A. Mangi, M. S. Senin, S. S. M. Zuki, M. H. W. Ibrahim  
510 - 526

[Effects of particle size and composition of cassava peels and rice husk on the briquette performance](#)

S. Anggraeni, S. N. Hofifah, A. B. D. Nandiyanto, M. R. Bilad  
527 - 542

[Analytical solution of the problem of heat exchange of buried pipeline with atmosphere in the conditions of snow cover](#)

E. V. Markov, S. A. Pulnikov, Y. S. Sysoev, D. V. Rachkov, E. P. Bragar

543 - 555

[Studying the effect of increasing capacity using comp technology in LTE-A networks](#)

A. N. Hammodat, S. A. Ayooob

556 - 570

[3D biomechanical analysis of swimming start movements using a portable smart platform with android pie](#)

A. Rusdiana, B. Mulyana, D. R. Nurjaya, Badruzaman, I. Imanudin, E. Fauziah, A. M. Syahid

571 - 585

[An Experimental identification of shear wave velocity of Bangkok clay using bender element](#)

S. Pethrung, P. Pairojn, S Teachavorasinskun

586 - 600

[Teaching the corrosion of iron particles in saline water to students with special needs](#)

R. Maryanti, A. Hufad, A. B. D. Nandiyanto, S. Tukimin

601 - 611

[Incineration alternative for municipal solid waste disposal of Najaf city, Iraq](#)

H M. M. Al-Khateeb

612 - 620

[Hybrid adjustment delay and superframe duration scheme to improve performance IEEE 802.15.4 networks](#)

E. Yundra, U. T. Kartini, I. W. Susila

621 - 634

[Students' perception of online learning and institution's engagement during the Covid-19 pandemic: A survey study of Taylor's design, architecture, and engineering students](#)

P. C. L. Koh, S. K. Wong

635 - 651

[A smartphone indoor localization based on affinity propagation clustering and Kullback-Leibler multivariate Gaussian](#)

O. A. Abdullah

652 - 665

[Digital literacy skills of vocational school teachers](#)

S. Saripudin, I. B. Budiyo, R. Listiana, A. Ana

666 - 680

[Adaptive cruise control of a simscape driveline vehicle model using PID controller](#)

A. Mahmood, M. Almagd, M. N. Noaman, Y. H. S. Alnema

681 - 695

[Low sulphidation Au-Ag epithermal associated minerals using T<sup>2</sup> hotelling's method](#)

D. N. Usman, L. Pulungan, S. Widayati, F. Hirnawan, N. Sulaksana, I. Haryanto, H. Nuryahya

696 - 710

[Deep-intrusion detection system with enhanced UNSW-NB15 dataset based on deep learning techniques](#)

A. M. Aleesa, M. Younis, A. A. Mohammed, N. M. Sahar

711 - 727

[E-monitoring the vulnerability of malnutrition of children using the TOPSIS method](#)

S. Nurhayati, R. Lubis

728 - 736

[Compare arithmetic coding to the wavelet approaches for medical image compression](#)

N. H. Salman

737 - 749

[Finite element analysis of two-way hollow high strength self-compacting fiber reinforced concrete slab under the effects of uniform repeated load \(using two-way piping system\)](#)

W. A. Tameemi, N. A. Alwash  
750 - 761

[Experimental investigation on an elevated basin solar still with integrated internal reflectors and inclined fins](#)

W. H. Alawee, H. A. Dhahad, I. S. Ahmed, T. A. Mohammad  
762 - 777

[Design and development of animation media for isomorphous binary phase diagram of engineering material courses](#)

M. Komaro, Ariyano, A. Suherman, A. Ana, R. Pangestu  
778 - 791

[Effect of curing time on compressive strength of agrostone building material with water hyacinth, bagasse and grass bio-fillers – a comparative study](#)

Y. M. Degu  
792 - 806

[Design and implementation of electronic examination device for improving the blind students' comfort](#)

H. Hidayat, N. S. Hasibuan, M. F. Wicaksono  
807 - 815

[Palmpoint recognition system based on proposed features extraction and \(c5.0\) decision tree, k-nearest neighbor \(KNN\) classification approaches](#)

M. S. Kadhm, H. Ayad, M. J. Mohammed  
816 - 831

[Prediction of the maximum throat width due to contraction in rectangular channels under critical flow conditions using computational fluid dynamics](#)

F. A. Alshitali, L. A. Alasadi, A. A. W. Alkraid, M. A. Aljumaili  
832 - 841

[An empirical numerical relation between evaporation rate and flash point of common industrial liquid materials](#)

J. F. Mata-Segreda  
842 - 850

[E-Rubric to measure employability skills](#)

S. Subekti, A. Ana, M. Muktiarni, V. Dwiyantri  
851 - 860

[Pneumonia detection using CNN through chest X-ray](#)

H. GM, M. K. Gourisaria, S. S. Rautaray, M. Pandey  
861 - 876

[Pelesir: Digital platform with a combination of frameworks to introduce tourism in southern of West Java, Indonesia](#)

G. Undang, A. M. Tanjung, T. Maskun  
877 - 887



## Editorial Board

[Home](#)

[Editorial Board](#)

[Submit a paper](#)

[Indexing and Awards](#)

[Reviewers](#)

[Articles in Press](#)

[Publication Ethics](#)

[Archives](#)

### Editor-In-Chief

- **Abdulkareem Shafiq Mahdi Al-Obaidi**, Ph.D.  
Associate Professor, School of Computer Science and Engineering  
Faculty of Innovation and Technology  
Taylor's University  
Taylor's Lakeside Campus  
No. 1 Jalan Taylor's, 47500 Subang Jaya  
Selangor DE  
Malaysia

### Editors

- **G. Davies**, Ph.D.  
Professor, Dean, Faculty School of Engineering  
The University of New South Wales  
UNSW Sydney  
NSW 2052  
Australia
- **Rodney Chaplin**, Ph.D.  
Associate, Professor, Associate Dean (International)  
Faculty of Engineering  
The University of New South Wales  
UNSW Sydney  
NSW 2052  
Australia
- **Andrew Ooi**, Ph.D.  
Associate, Professor, Assistant Dean (International)  
School of Engineering  
The University of Melbourne  
Victoria 3010  
Australia
- **David WL Hukins**, Ph.D.  
B.Sc., Ph.D. (London), D.Sc. (Manchester), C.Phys., F.Inst.P., F.I.P.E.M., F.R.S.E.  
Professor of Bio-medical Engineering  
Head of Mechanical and Manufacturing Engineering  
School of Engineering  
Mechanical Engineering  
The University of Birmingham  
Edgbaston  
Birmingham  
B15 2TT  
United Kingdom
- **Takayuki Saito**, Ph.D.  
Professor, Shizuoka University  
Graduate School of Science and Engineering  
3-5-1 Johoku  
Hamamatsu  
Shizuoka 432-8561  
Japan
- **S. B. Chin**, Ph.D.  
Professor, The University of Sheffield  
Mechanical Engineering Department  
Mappin Street, Sheffield  
S1 3JD,  
United Kingdom
- **Xiaoyu Luo**, Ph.D.  
Professor, Department of Mathematics  
University of Glasgow  
Glasgow G12 8QW
- **Stephen B M Beck**, Ph.D.  
Professor in Mechanical Engineering  
Faculty Director of Learning and Teaching – Engineering  
Department of Mechanical Engineering

The University of Sheffield  
Mappin Street  
Sheffield  
S1 3JD  
United Kingdom

- **Xiao (Yun) Xu**, Ph.D.  
Professor of Biofluid Mechanics  
Department of Chemical Engineering  
Imperial College  
London  
United Kingdom
- **Seeram Ramakrishna**, Ph.D.  
Professor, Dean, Faculty of Engineering  
Dean's Office, Block EA, #07-26  
9 Engineering Drive 1,  
National University of Singapore,  
Singapore 117576
- **Ramesh Singh Kuldip Singh**, Ph.D.  
Senior Professor of Mechanical and Materials Engineering  
Faculty of Engineering  
University of Malaya, 50603 Kuala Lumpur, Malaysia  
Universiti Teknologi Brunei, BE1410 Gadong, Brunei Darussalam
- **Gary Hawley**, Ph.D.  
Professor of Automotive Engineering  
Dean and Medlock Chair of Engineering,  
Faculty of Engineering and Design  
University of Bath,  
Claverton Down, Bath BA2 7AY  
United Kingdom
- **Yousif Abdall Abakr**, Ph.D.  
School of Mechanical Engineering  
The University of Nottingham, Malaysia Campus  
Jalan Broga, 43500 Semenyih, Selangor  
Malaysia
- **R. Rajesh @ Nithyanandam**, Ph.D, PGChETL, MICheM  
Professor, Department of Chemical Engineering  
Mohamed Sathak Engineering College  
Kilakarai, Tamil Nadu  
(Affiliated to Anna University)  
India
- **Chong Perk Lin**, Ph.D, PgCLTHE, FHEA, CEng, MIMechE  
Senior Lecturer of Mechanical Engineering  
Programme Leader of BEng Tech (Hons) Mechanical Engineering  
School of Computing, Engineering and Digital Technologies  
Teesside University, Middlesbrough, Tees Valley. TS1 3BX.  
United Kingdom.



## VETIVER WASTE FERTILIZER AS A GROWTH CATALYST ON MUNG BEAN

EDDY SOERYANTO SOEGOTO<sup>1</sup>,  
RIZKY JUMANSYAH<sup>2,\*</sup>, SENNY LUCKYARDI<sup>3</sup>

<sup>1</sup>Departemen Manajemen, Universitas Komputer Indonesia, Indonesia

<sup>2\*</sup>Departemen Sistem Informasi, Universitas Komputer Indonesia, Indonesia

<sup>3</sup>Departemen Magister Manajemen, Universitas Komputer Indonesia, Indonesia

\*Corresponding Author: jumarizky@mahasiswa.unikom.ac.id

### Abstract

Mung bean (*Vigna radiata*) is extensively cultivated in tropical and sub-tropical Asia and a resource of seed protein. It is also easily adapted to various environments. Fertilizer has an important role in increasing production and quality of crop cultivation. The fertilizers must meet national standards and their effectiveness must be proven. The purpose of this study is to analyse the use of vetiver waste fertilizer as growth catalyst on mung bean. This research method used completely randomized design (CRD) and descriptive qualitative method. This research was conducted by observing three types of sample, namely the mung bean without fertilizer, half-decomposed fertilizer, and decomposed vetiver waste fertilizer. The sample of mung bean without fertilizer was used as a reference for standardization of mung bean growth. In addition, this study discusses the difference of using vetiver waste fertilizer on plant physical characteristics. The results showed that from the samples taken, mung bean with half-decomposed fertilizer were more stunted than the other two samples. Meanwhile, from the calculation using CRD is obtained that value F calculate < F tables. It means that even though special treatment is given, it does not have significant effect on the Mung bean growth due to it will continue to grow even if there is no fertilizer application. However, plants with decomposed vetiver waste fertilizer have rapid growth with stronger plant physical characteristics. It is indicated by more water contained in the mung bean with decomposed vetiver waste fertilizer. From these results, it is shown that the use of vetiver waste fertilizer is effective as catalyst in the growth of mung bean. This research is expected to be a reference for future research. It can be concluded that vetiver waste fertilizer can stimulate the growth of mung bean and potentially effective for the development of other plants.

Keywords: Agricultural, Catalyst, Mung bean, Vetiver waste fertilizer.

## 1. Introduction

Fertilizer has an important role in increasing the production and quality of crop cultivation. To meet quality standards and ensure the effectiveness of fertilizers, the fertilizers produced must be tested. Fertilizers can be classified into several categories based on origin, chemical, and physical characteristics. According to its origin, fertilizers can be divided into natural and artificial fertilizers. Based on the chemicals, it can be classified into organic and inorganic. From its physical characteristics, it can be divided into solid and liquid fertilizer. Generally, fertilizers that are often used by farmers are urea, ZA (Zwavelzure Amonim), SP36 (Super Phospate), KCI (Potassium Chloride), ZK (Zwalvera Kali), NPK PHONSKA (Nitrogen Phospate Potassium), and Dolomite (Lime Carbonate) [1, 2]. However, the use of urea is minimized and started to be replaced by organic fertilizers. It is because organic fertilizers can affect the physical characteristics and chemical properties of the soil, such as the stability of aggregates and the ability to hold groundwater so that it can maintain water needs for plants.

The increase in soil aggregate stability by organic fertilizers is caused by the presence of gum polysaccharides obtained from soil bacteria, the presence of hyphae and fungal growth from actinomycetes around soil particles [3, 4]. Besides, organic fertilizers also play a role in increasing physical, chemical, and biological fertility of the soil so that it can improve the efficiency of inorganic fertilizers such as urea. The quality and composition of organic fertilizers varies depending on the compost base material and the manufacturing process. The use of legume plants in the form of alley cropping and cover crops as well as in-situ organic matter needs to be intensified to support the use of non-commercial organic fertilizers and restore soil fertility [5].

Vetiver grass are plants that require constant light and are able to grow on land that has a very wide pH range from 3.5 (very acidic) to 11.5 (very alkaline). In order to produce high oil content and quality, vetiver should be planted in sandy soil with an altitude of more than 750 m ASL [6]. Previous research on vetiver plants showed that this plant functions as soil conservation because it can reduce the level of soil erosion and heavy metals contamination such as Zn, Cu, and Fe [7, 8]. In addition, other studies also show that vetiver plants can also be used as compost and raw material for paper and handicraft industries, while the roots are used for woven crafts and insect repellent [9]. As for further research, it is found that this plant has an economic value that can contribute to the country's foreign exchange products from its oil, namely vetiver oil. Essential oil produced from the roots of vetiver plant is widely used as a binder in perfume production because of its strong fixing power as well as raw material for cosmetics and medicines [10].

Meanwhile, a research conducted by Emenecker and Strader related to plant growth and development showed the influence of plant hormones. Plant hormones play a role in regulating the growth, development, and response of plants to biotic and abiotic stress. The types of plant hormones that play a comprehensive role in plant growth and development (especially in the stages of root lengthening and seed germination) are Auxin and Abscisic Acid (ABA) [11]. Moreover, the research on growth inhibition in plants shows that one of the things that inhibits plant growth and development is the content of lead (Pb) in the soil. For plant types such as mung bean, the inhibiting factors for seed development and growth are influenced by saline oil, so that Plant Growth Promoting Rhizobacteria (PGPR) and Silicon (Si)

are needed to reduce the effects of salinity stress [12]. The factors that affect the quality of mung beans are Relative Humidity (RH), light, and Plasma-Activated Water (PAW), this is because the higher the RH value indicates higher water absorption and conductivity, compared to the lower RH value. Induction of light helps increase plant water transport and antioxidant production in mung bean growth. Also, PAW affects the growth of mung beans because it has active components in PAW, such as NO-2 -, NO-3, and H2O2 [13, 14].

## 2. Research Method

This study used the CRD method with the following formula:

Fully Randomized Design (FRD) [15]:-

$$t(n - 1) \geq 15 \tag{1}$$

$$FK: \frac{y...^2}{nxt} \tag{2}$$

$$JKT : \sum_{i=1}^t \sum_{j=1}^n Yij^2 - FK \tag{3}$$

$$JKP : \sum_{t=1}^t \frac{Yi^2}{n} - FK \tag{4}$$

$$JKG : JKT - JKP \tag{5}$$

$$KTP : \frac{JKP}{t-1} \tag{6}$$

$$KTG : \frac{JKG}{t(n-1)} \tag{7}$$

$$F \text{ Count} : \frac{KTP}{KTG} \tag{8}$$

Therefore, the various effects of treatment on green bean plants are the following (Table 1):

**Table 1. The various effects of treatment on green bean plants**

Source of diversity (S.K)	Degree of freedom (d.b)	Sum of squares (J.k)	Middle square (K.T)	F Count	F table
					0.05 0.01
<b>Treatment</b>	<i>t</i> -1	JKP	KTP		
<b>Attempt Error</b>	<i>t</i> ( <i>n</i> -1)	JKG	KTG		
<b>Total</b>	<i>tn</i> -1	JKT			

where: *t*: Much Treatment, *n*: Many Repetitions, *i*: Number of Treatments, *j*: Number of Observations, FK: Correction Factors, JKT: Total Sum of Squares, JKP: Sum of Squares between Treatments, JKG: Sum of the Squares of the Error, KTP: Middle Square Treatment, KTG: Middle Square Error, F Count: F Value of Analysis Calculation Results

This study used a completely randomized design (CRD) and descriptive qualitative method. CRD is the simplest design, suitable for laboratory research, greenhouses or experiments that have homogeneous experimental materials. The advantage of this method is that the experimental design plan is relatively easy, statistical analysis is very simple, and it is flexible in the number of treatments or repetitions [15]. CRD is applied in this study because the material is relatively homogeneous, namely using cotton as a growing medium and a limited number of

treatments. We conducted direct observations on the growth of mung bean stem for five days indoor. Mung bean seeds were placed in three Petri dishes with different treatments. The first treatment is conducted without using vetiver waste fertilizer. The second treatment used half-decomposed vetiver waste fertilizer. The third treatment used vetiver waste fertilizer that had been decomposed. Planting mung bean seeds using cotton media with a weight of 2 grams in each petri dish. For the second and third treatment using fertilizer weighing 1 gram measured using Nankai ART: 177-21 scales. The fertilizer is grinded using a Miyako blender (3 in 1) BL - 102 PL, with room temperature ranging from 19 to 20°C and humidity ranging from 62 to 65%. Stem measurements were carried out using the Microscope Micrometer Calibration Ruler.

### 3. Results and Discussion

The use of fertilizer is one of the most important factors to increase plant productivity [16]. Vetiver waste is used as an organic fertilizer for plants. It is shown by the results of this study. In the first treatment, which was carried out without applying vetiver waste fertilizer, the growth of mung beans was normal. It is because the main food source is in the seeds (endosperm) and can be obtained through water. The second and third treatments with vetiver waste fertilizer showed different results. The second treatment used half-decomposed vetiver waste fertilizer showed the growth of mung bean seeds was inhibited. The third treatment with decomposed vetiver waste fertilizer showed the fastest growth.

The stem length in the first and second treatment was 40 and 18 mm. Meanwhile, the third treatment was 45 mm. Observation on stem water content showed that the third treatment had more water content than the first and second treatments. However, the third treatment had a faster leaf growth than the first and second treatments. In addition, observations on the roots showed that the roots of mung beans in the third treatment were stronger than the first and second treatments.

This study captures that fertilization using vetiver waste can increase plant growth [17, 18]. Nutrition at the growth stage is very important for mung bean to improve the plant performance in terms of growth, yield, and quality of mung bean (Krishnaveni et al. 2004). Seeds still require low amounts of nutrients because they are able to absorb their needs from the cotyledons.

Factors that influence the growth of mung beans are light intensity, nutrients, water, humidity, and temperature [19]. Nutrients is obtained through fertilization. Table 2 shows the growth data of mung beans without using vetiver waste fertilizer.

The first day of planting without fertilizer did not show any changes in plant length with a temperature of 19.7°C and humidity of 65%. On the second day, there was a change in temperature which increased to 20.1°C but there was no change in humidity, which was still 65%. Changes occurred in plant length, where the first day the plant length was 0 mm but on the second day the plant length became 6 mm. On the third day, the room temperature and humidity decreased to 19.2°C and 62%. However, the growth in plants continued to increase. The length of the plant which was previously only 6 mm increased to 15 mm. On the fourth day the temperature remained stable, namely 19.2°C, with a humidity that increased by 1%. The length of the plant was enlarged by 12 mm, where the plant size was 27 mm.

On the fifth day, the temperature and humidity increased to 20.1°C and 64%. On the fifth day, the length of the plant increased to 40 mm.

From the table 2, based on the results of mung bean growth showed that the growth of mung bean plants without fertilizer can grow normally. Meanwhile, the plant growth without fertilizers was better than the half-decomposed fertilizers. In addition, it differs only a few millimetres in size from plants that use decomposed fertilizers. With this comparison, it can be a consideration for the farmers to use vetiver waste fertilizers (6 months of decomposition). From the data shown in Table 2, it can be different from the results of other studies or even in its application. It is because several factors such as light intensity, temperature, and water availability in plants.

**Table 2. Sample growth data without fertilizer.**

<b>Days to</b>	<b>Temperature</b>	<b>Humidity</b>	<b>Plant Length</b>
<b>Day 1</b>	19.7°C	65%	0 mm
<b>Day 2</b>	20.1 °C	65%	6 mm
<b>Day 3</b>	19.2 °C	62%	15 mm
<b>Day 4</b>	19.2 °C	63 %	27 mm
<b>Day 5</b>	20.1 °C	64%	40 mm

In this study, in general the three plants with different treatments were still able to grow. It is because these plants are treated indoors with stable temperatures and humidity. Suitable environmental conditions are always needed for plants to grow. However, the environmental conditions in which the plants are located are always changing. If the plant can still grow, the environmental changes that occur are still within the tolerance limit of the plant. However, plants often experience extreme environmental changes that can cause decreased productivity and even death. It shows that each plant has a limiting factor and tolerance to the environment [20, 21].

This research was conducted with the same light treatment. It causes plant growth to remain normal without being affected by lighting constraints. Because plants are placed without direct sunlight, they grow quickly and avoid etiolation. Etiolation is plants that are not exposed to light, so auxin stimulates the elongation of cells so that they grow longer, but are less healthy, thin, and pale. On the other hand, a lot of auxin light is damaged so that plant growth is inhibited. The broken auxin that is dispersed to the darker side of the plant is caused by a lack of light. The growth of the stems decreases and becomes shorter. However, the plants are sturdier, the leaves are fully grown, and greener. The process of photosynthesis in plants requires light for the formation of chlorophyll. Contrariwise, chlorophyll is damaged if the light intensity is too high [22]. The role of fertilizers in the effectiveness of plant fertility is quite significant. The addition of fertilizers plays a role in improving the physical characteristics of the soil by reducing density and binding water for plant growth needs [23]. In this study, vetiver waste fertilizers were classified into two types, namely half-decomposed (decomposed for 3 months) and decomposed fertilizers. We analysed a comparative impact of the two fertilizers. The following is a display of half-decomposed fertilizers (see Fig. 1).

The positive effect of fertilizers is to stimulate the growth of sprouts [24]. In this study, we analysed the growth process of mung bean with half-decomposed

vetiver waste fertilizer. Organic fertilizers have a significant impact to increase growth (See Table 3).



**Fig. 1. Half-decomposed vetiver waste fertilizer.**

**Table 3. Sample growth data with half-decomposed.**

<b>Days</b>	<b>Temperature</b>	<b>Humidity</b>	<b>Plant Length</b>
<b>Day 1</b>	19.7 °C	65%	0 mm
<b>Day 2</b>	20.1 °C	65%	3 mm
<b>Day 3</b>	19.2 °C	62%	11 mm
<b>Day 4</b>	19.2 °C	63 %	13 mm
<b>Day 5</b>	20.1 °C	64%	18 mm

In the first day, there was no growth occurred in the temperature and humidity of 19.7°C and 65%. In the second day, a growth of 3 mm occurred in the temperature of 20.1°C and humidity of 65%. In the third day, the growth continued to increase to 11 mm. The temperature and humidity were 19.2°C and 62%. In the fourth day, the length reached 13 mm. Therefore, the temperature and humidity in the fourth day were 19.2°C and 63%. After five days of observing mung beans with half-decomposed vetiver waste fertilizer, the final result showed 18 mm in length with average growth of 9 mm. The growth was in the average temperature of 19.6°C and humidity of 63.8%.

Compared to the growth of mung beans without using fertilizer and decomposed fertilizer, the mung beans growth with half-decomposed fertilizer is the slowest grow. It happened because half-decomposed organic fertilizer did not work optimally. Fertilizer that is intended to increase the crop yields shows the opposite because of the nutrients in half-decomposed organic fertilizer [25].

The addition of fertilizer such as bio-fertilizer can affect plant growth [26]. In addition, we conducted this research by observing the growth of mung bean by using fertilizer. The test was conducted within five days by measuring the air temperature, humidity, and the length of the plant. In this research, we used the Air Quality Monitor to measure the temperature and humidity. The results of the test are shown in Table 4.

The test was conducted in five days and consecutively checked every 10.23 AM to see the growth of the mung beans, temperature, and humidity. On the first day,

the plant length was 0 mm with air temperature of 19.7°C and humidity of 65%. The second day, the length of the plant grew by 7 mm while humidity remains the same as the first day. However, the temperature changed and increased by 0.4°C, which made it changed from 19.7°C to 20.1°C. On the third day, the plant length grew by 16 mm, which made it became 23 mm. Nevertheless, the temperature and humidity had decreased significantly by 0.9°C and 3% which made it became 19.2°C and 62%. On the fourth day, the length of the plant grew by 8 mm, which made it became 31 mm with the same temperature as the third day. However, the humidity increased by 1%, which made it became 63%. On the last day, the final plant length grew by 14 mm, which made it became 45 mm. The air temperature on the last day increased significantly by 0.9°C and made it changed to 20.1°C that was same with the temperature on the second day. The humidity was increased by 1%, which made it became 64%. As seen on the Table 4, the usage of vetiver waste fertilizer shows a significant increased growth in the mung bean than the half-decomposed vetiver waste fertilizer and without using fertilizer. The mung bean sprout can grow up to 45 mm in five days with the help of this bio-fertilizer. It is in-line with the results of Mishra et al. that stated bio-fertilizers is an essential supporting component that play vital role in plant growth [27].

**Table 4. The results of mung bean growth with fertilizer.**

Day	Temperature	Humidity	Plant Length
Day 1	19.7°C	65%	0 mm
Day 2	20.1°C	65%	7 mm
Day 3	19.2°C	62%	23 mm
Day 4	19.2°C	63%	31 mm
Day 5	20.1°C	64%	45 mm

Mung beans contain protein isolate (MuPI) with a concentration of 8S globulin (80%), which shows homology (68%) and structural b-conglycinin. This content provides benefits to ward off free radicals and fight cancer cell lines [28, 29]. Meanwhile, during the mung bean sprouts growth, its seeds utilize energy sources obtained from water and food reserves stored in the endosperm to produce a higher succinic acid content [30]. Then, without using fertilizer when the mung bean seeds grow into sprouts, the external factor that affects their growth is water. The more water absorbed by the roots, the faster growth of mung bean stem. It can also affect the size of the stem. Stem that grows without using vetiver waste fertilizer and using vetiver waste fertilizer have different water content. It is due to the texture of fertilizers that have decomposed completely and half-decomposed have different textures and shapes. Fertilizer that has decomposed completely has a smoother texture. Therefore, stem without vetiver waste fertilizer have sufficient water content in the stem. Meanwhile, stem with vetiver waste fertilizers that have undergone a decomposition process have more water content in the stems. On the other hand, stem that use vetiver waste that has not gone through a complete decomposition process to become fertilizer have less water content in the stems (Fig. 2). It affects the quality and size growth of mung bean stem.

Vetiver waste fertilizer that has undergone a decomposition process helps the stem to maintain moisture and water availability. Also, it reduces the duration of evaporation of water into the air due to sunlight. On the other hand, vetiver waste that has not undergone a perfect decomposition process as fertilizer inhibits the

absorption of water needed by the stem. It is because the vetiver waste still has the property of absorbing water and has a salt content. Therefore, the water that should be absorbed by the stem is locked in the half-decomposed vetiver waste fertilizer [31, 32].




Fig. 2. Comparison of stems of mung bean stem.

### Complete randomized design of mung bean plants

This study used a completely randomized design method to determine the significance of the effect of using vetiver fertilizers on the growth of green beans, with the following calculations:

Fully Randomized Design (FRD)

$$4(n - 1) \geq 15$$

$$n = \frac{11}{4} = 2.75 = 3$$

So at least a lot of repetition of green bean growth observations that must be done is as many as 3 experiments

The value of the Correction Factor (FK) carried out is as follows:

$$FK: \frac{y \dots^2}{nxt} = \frac{(137)^2}{12} = 1564.083$$

Meanwhile, to find out the Sum of Squares of Errors (JKG), first, you must calculate the value of the Total Squares (JKT) and the Sum of Squares between Treatments (JKP) as follows:

$$\begin{aligned} JKT &= (0)^2 + (15.5)^2 + (0)^2 + (0)^2 + (38)^2 + (10.5)^2 + (3)^2 + (15)^2 + (7)^2 \\ &\quad + (0)^2 + (35.5)^2 + (11.5)^2 - FK \\ &= 3470 - 1564.083 \\ &= 1905.917 \end{aligned}$$

$$\begin{aligned} JKP &= (45)^2 + (26)^2 + (38.5)^2 + (27.5)^2 - FK \\ &= \frac{4939.5}{3} - 1564.083 \\ &= 82.417 \end{aligned}$$

$$\begin{aligned} JKG &= 1905.917 - 82.417 \\ &= 1823.5 \end{aligned}$$



Then the Middle Square (KT) calculation stage is carried out and consists of two types, namely the Treatment Middle Square (KTP) and the Error Middle Square (KTG) as follows:

$$\begin{aligned} \text{Middle Square:} \\ \text{KTP} &= \frac{82.413}{3} = 27.472 \\ \text{KTG} &= \frac{1823.5}{8} = 227.9375 \end{aligned}$$

The F value of the Analysis Calculation Results is as follows:

$$F \text{ Count} : \frac{\text{KTP}}{\text{KTG}} = \frac{27.472}{227.9375} = 0.1205$$

Therefore, the analysis of the various treatment effects on the growth of green bean plants obtained the value of F count <F table namely 0.1205 < 4.07 and 7.59. Thus, it was found that different treatments for seeds such as fertilizer application did not significantly affect the growth of mung bean plants. This is because there are other factors that affect plant growth namely water, air, soil, humidity, light, and maintenance.

#### 4. Conclusion

From the results of this study, it can be concluded that the use of vetiver waste fertilizer on mung bean plants is considered effective. It is supported by the results of the differences between plants that use half-decomposed and decomposed vetiver waste fertilizer. Based on the reference for plants that do not use fertilizers, the difference in stem length is 5 mm and 27 mm, with an average temperature and humidity of 19.6°C and 63.8%. The physical condition of plants that use vetiver waste fertilizer has more nutrient availability so that growth can be faster with maximum nutrient absorption. Also, this study also identified that not all plants that were given special treatment (applying fertilizers) would experience significant changes in the plant's growth.

#### References

1. Firmansyah, I.; Spiller, M.; De Ruijter, F.J.; Carsjens, G.J.; and Zeeman, G. (2017). Assessment of nitrogen and phosphorus flows in agricultural and urban systems in a small island under limited data availability. *Science of the Total Environment*, 574, 1521-1532.
2. Mahyudin, R.P.; Firmansyah, M.; Anggraini, M.; and Najmina, D. (2020). Bioremediation of iron on diamond post mining soil using compost made from cow manure and traditional market organic waste. *Journal of Ecological Engineering*, 21(5), 221-228.
3. Kim, S.; Simpson, A.J.; Kujawinski, E.B.; Freitas, M.A.; and Hatcher, P.G. (2003). High resolution electrospray ionization mass spectrometry and 2D solution NMR for the analysis of DOM extracted by C18 solid phase disk. *Organic Geochemistry*, 34(9), 1325-1335.
4. Dong, D.; Nelson, Y.M.; Lion, L.W.; Shuler, M.L.; and Ghiorse, W.C. (2000). Adsorption of Pb and Cd onto metal oxides and organic material in natural surface coatings as determined by selective extractions: new evidence for the importance of Mn and Fe oxides. *Water Research*, 34(2), 427-436.

5. Gao, C.; El-Sawah, A.M.; Ali, D.F.I.; Hamoud, Y.A.; Shaghaleh, H.; and Sheteiw, M.S. (2020). The integration of bio and organic fertilizers improve plant growth, grain yield, quality, and metabolism of hybrid maize (*Zea mays* L.). *Agronomy*, 10(3), 319.
6. Aishwath, O.P.; and Ratan, L. (2016). Resilience of spices, medicinal and aromatic plants with climate change induced abiotic stresses. *Annals of Plant and Soil Research*, 18(2), 91-109.
7. Singh, S.; Fulzele, D.P.; and Kaushik, C.P. (2016). Potential of vetiveria zizanioides L. Nash for phytoremediation of plutonium (<sup>239</sup>Pu): chelate assisted uptake and translocation. *Ecotoxicology and Environmental Safety*, 132, 140-144.
8. Wilde, E.W.; Brigmon, R.L.; Dunn, D.L.; Heitkamp, M.A.; and Dagnan, D.C. (2005). Phytoextraction of lead from firing range soil by vetiver grass. *Chemosphere*, 61(10), 1451-1457.
9. Liu, H.; Xie, Y.; Li, J.; Zeng, G.; Li, H.; Xu, F.; Feng, S.; and Xu, H. (2020). Effect of serratia sp. K3 combined with organic materials on cadmium migration in soil-vetiveria zizanioides L. system and bacterial community in contaminated soil. *Chemosphere*, 242, 125164.
10. David, A.; Wang, F.; Sun, X.; Li, H.; Lin, J.; Li, P.; and Deng, G. (2019). Chemical composition, antioxidant, and antimicrobial activities of vetiveria zizanioides (L.) nash essential oil extracted by carbon dioxide expanded ethanol. *Molecules*, 24(10), 1897.
11. Emenecker, R.J.; and Strader, L.C. (2020). Auxin-abscisic acid interactions in plant growth and development. *Biomolecules*, 10(2), 281.
12. Ahmed, N.; Ahsen, S.; Ali, M.A.; Hussain, M.B.; Hussain, S.B.; Rasheed, M.K.; Butt, B.; Irshad, I.; and Danish, S. (2020). Rhizobacteria and silicon synergy modulates the growth, nutrition and yield of mungbean under saline soil. *Pakistan Journal Botany*, 52(1), 9-15.
13. Amitrano, C.; Arena, C.; De Pascale, S.; and De Micco, V. (2020). Light and low relative humidity increase antioxidants content in mung bean (*Vigna radiata* L.) sprouts. *Plants*, 9(9), 1093.
14. Fan, L.; Liu, X.; Ma, Y.; and Xiang, Q. (2020). Effects of plasma-activated water treatment on seed germination and growth of mung bean sprouts. *Journal of Taibah University for Science*, 14(1), 823-830.
15. Mattjik, A.A.; and Sumertajaya, I.M. (2013). Perancangan percobaan dengan aplikasi sas dan minitab. *PT Penerbit IPB Press*.
16. Godfray, H.C.J.; Beddington, J.R.; Crute, I.R.; Haddad, L.; Lawrence, D.; Muir, J.F.; Pretty, J.; Robinson, S.; Thomas, S.M. and Toulmin, C. (2010). Food security: the challenge of feeding 9 billion people. *Science*, 327(5967), 812-818.
17. Fernández, V.; and Brown, P.H. (2013). From plant surface to plant metabolism: the uncertain fate of foliar-applied nutrients. *Frontiers in Plant Science*, 4, 1-15.
18. Krishnaveni, S.; Anandha, A.; Palchamy; and S. Mahendran. (2004). Effect of foliar spray of nutrients on growth and yield of green gram (*Phaseolus radiatus*). *Legume Research-An International Journal*, 27(2), 149-150.
19. Srivastava, G.; and Mehrotra, R.C. (2010). New legume fruits from the oligocene sediments of assam. *Journal of the Geological Society of India*, 75(6), 820-828.

20. Pandey, N.; and Bhatt, R. (2016). Role of soil associated exiguobacterium in reducing arsenic toxicity and promoting plant growth in vigna radiata. *European Journal of Soil Biology*, 75, 142-150.
21. Castanha, N.; Miano, A.C.; Sabadoti, V.D.; and Augusto, P.E.D. (2019). Irradiation of mung beans (*Vigna radiata*): a prospective study correlating the properties of starch and grains. *International Journal of Biological Macromolecules*, 129, 460-470.
22. Sarwar, R.A.; Ashraf, M.; and Javaid, M.M. (2018). Response of foliar-applied nutrient solution with and without soil-applied fertilizers on growth and yield of mung bean. *Journal of Plant Nutrition*, 41(9), 1083-1093.
23. Weber, J.; Karczewska, A.; Drozd, J.; Licznar, M.; Licznar, S.; Jamroz, E. and Kocowicz, A. (2007). Agricultural and ecological aspects of a sandy soil as affected by the application of municipal solid waste composts. *Soil Biology and Biochemistry*, 39, 1294-1302.
24. El-Din, S.M.M. (2015). Utilization of seaweed extracts as bio-fertilizers to stimulate the growth of wheat seedlings. *The Egyptian Journal of Experimental Biology*, 11, 31-39.
25. Haynes, R.J.; and Naidu, R. (1998). Influence of lime, fertilizer and manure applications on soil organic matter content and soil physical conditions: a review. *Nutrient Cycling in Agroecosystems*, 51(2), 123-137.
26. Khalilzadeh, R.; Tajbakhsh, M.; and Jalilian, J. (2012). Growth characteristics of mung bean (*Vigna radiata* L.) affected by foliar application of urea and bio-organic fertilizers. *International Journal of Agriculture and Crop Sciences (IJACS)*, 4(10), 637-642.
27. Mishra, D.; Rajvir, S.; Mishra, U.; and Kumar, S.S. (2013). Role of bio-fertilizer in organic agriculture: a review. *Research Journal of Recent Sciences* 2, 39-41. (ISSN), 2277, 2502.
28. Watanabe, H.; Inaba, Y.; Kimura, K.; Asahara, S.I.; Kido, Y.; Matsumoto, M.; Tachibana, N.; Kaneko, S.; Kohno, M.; and Inoue, H. (2017). Dietary mung bean protein reduces hepatic steatosis, fibrosis, and inflammation in male mice with diet-induced, nonalcoholic fatty liver disease. *The Journal of Nutrition*, 147(1), 52-60.
29. Gupta, N.; Srivastava, N.; and Bhagyawant, S.S. (2018). Vicilin - A major storage protein of mungbean exhibits antioxidative potential, antiproliferative effects and ACE inhibitory activity. *PLoS One*, 13(2), e0191265.
30. Chen, L.; Wu, J.E.; Li, Z.; Liu, Q.; Zhao, X.; and Yang, H. (2019). Metabolomic analysis of energy regulated germination and sprouting of organic mung bean (*Vigna radiata*) using NMR spectroscopy. *Food chemistry*, 286, 87-97.
31. Islam, M.P.; Bhuiyan, M.K.H.; and Hossain, M.Z. (2008). Vetiver grass as a potential resource for rural development in Bangladesh. *Agricultural Engineering International: CIGR Journal*, 10(5), 1-18.
32. Cuong, D.C.; Van Minh, V.; and Truong, P. (2015). Effects of sea water salinity on the growth of vetiver grass (*Chrysopogon Zizanioides* L.). *Modern Environmental Science and Engineering*, 1, 185-191.